
The Progenitors of Dwarf Galaxies in Galaxy Clusters

> Steve Crawford @astrocrawford South African Astronomical Observatory

Collaborators: Matthew Bershady (U.Wisconsin), Greg Wirth (Keck Observatory), Solohery Randriamampandry (SAAO)


## Dwarf Ellipticals in clusters



Dwarf Ellipticals are heterogeneous class that is the most numerous in clusters

## But how did they get there?


# Build up of the faint end of the red sequence?



See also Capozzi, Collins & Stott 2010, Bildfell et al. 2012, De Lucia et al. 2007, Gilbank & Balogh 2008, Huertas-Company et al. 2009, Lemaux et al. 2012, Rudnick et al. 2012, Fassbender et al. 2014)

Also see Andreon (2008), Andreon et al. (2014), Lidman et al. (2008), and De Propris, Phillipps & Bremer (2013), Cerulo et al. (2017)

# Luminous Compact Blue Galaxies



- Original discovered by Koo & Kron in 80s as an observational class: unresolved blue galaxies
- Rapidly evolution *heterogeneous* population of galaxies (factor of ~10 drop since z~1, Guzman et al. 1997)
- Luminous ( $M_B \sim -20$ ), small ( $r_e \sim 2$  kpc), and intense star formation rates



HST/WFPC2/NICMOS

Ref: Koo et al. 1994, Koo et al. 1997, Guzman et al. 1996; Phillips et al. 1997; Kobulnicky & Zaritsky 1999; Guzma'n et al. 2003; Garland et al. 2004; Werk et al. 2004; Barton et al. 2006; Noeske et al. 2006; Rawat et al. 2007; Hoyos et al. 2007; Tollerud et al. 2010

## LCBG are like ...

**LOW-Z** <3.5 Gyrs ago z < 0.3

Blue Compact Dwarfs, HII galaxies



green peas




Cardamone et al. 2009

**Int-z** Up to 3.5-9 Gyrs ago 0.3 < z < 1

> Extreme Emission Line Galaxies


**CNELGs** 

High-z >9 Gyrs ago z>1 Lyman Break Galaxies



Burgarella et al. 2009

## LCBGs in Clusters



Handful of LCBGs in CL0024 seem to have similar properties to low redshift dwarf galaxies

Koo original proposed LCBGs as the progenitors of dE

Subset of the Butcher-Oemler Galaxies, but what are their properties and where do they go?


#### Observations

Field (1)	WLTV ID (2)	α (J2000) (3)	(J2000) (4)	z (5)	$(\mathrm{km \ s^{-1}})$ (6)	$R_{200}$ (Mpc) (7)	$R_{200}$ (") (8)
Cl 0016+16	w01	00:18:33.6	+16:26:16	0.5467	1490	2.74	428
Cl J1324+3011	w08	13:24:48.8	+30:11:39	0.7549	806	1.31	178
MS 1054-03	w07	10:56:60.0	-03:37:36	0.8307	1105	1.72	225
Cl J1604+4304	w10	16:04:24.0	+43:04:39	0.9005	1106	1.65	211

Table 1. Summary of Fields


Deep WIYN narrow band imaging combined with DEIMOS spectra, Archive HST, Spitzer, and VLA observations








## Identifying Cluster LCBGs




#### "Shell-like" LCBG Radial Distribution



Similar to results for low-z SF galaxies e.g.Thompson 1986; Ellingson et al. 2001; Mahajan et al. 2010


Crawford et al. 2011, 2014

# LCBGs are falling into the cluster



Crawford et al. 2014

## Young burst in dE




Rys et al. showed that the typical dE had a burst of star formation ~5 Gyrs ago

Also see Michielson et al 08, Lelli et al. 2014, Toloba et al. 2014, Mentz et al 2016

Rys et al. 2015


#### Bursts in Illustris Simulations

#### Dwarfs galaxies undergo a burst when entering the cluster



Mistani et al. 2015


### Spectroscopic Properties



Cluster and field LCBGs nearly indistinguishable in terms of dynamical mass, SFR, abundance, or size.


> Typical properties:  $\sigma \sim 56 \text{ km/s},$   $r_{1/2} \sim 1.8 \text{ kpc}$   $M_{dyn} \sim 5 \times 10^9 \text{ M}_{\odot}$  $12 + \log(O/H) = 8.6.$

### Spectroscopic Properties



Distribution of cluster dE

#### Dynamical to Stellar Mass




LCBG M<sub>dyn</sub>/M\* Cluster~ 2.6 Field ~ 4.8  $\frac{dE M_{dyn}/M*}{Cluster~ 2.2}$ Field ~ 5.1 Penny et al. 2015

Randriamapandry et al. 2017 <u>https://arxiv.org/abs/1706.04534</u>

## "Longitudinal Study"


#### **Observed Evolution**

#### Simulated Evolution




Mistani et al. 2015

# Clusters Triggering LCBGs




## Fate of LCBGs

30-75% of dE went through an LCBG phase between z=0.3-1




## **Complex Star formation in LCBGs**



Crawford et al. 2016

Range of star formation in different metrics

Starbursting galaxies need better modeling




Randriamampandry et al. 2015

# Summary


- Galaxy Clusters trigger the star burst phase in inflating dwarf galaxies at intermediate redshifts
- Spectral properties of LCBGs are very similar to local, cluster dE
- Likely between 30-75% of dE experienced a LCBG phase in the last 7.5 Gyrs
- Further work needed to study the evolution in dynamical to stellar mass, morphology/size, and complex star bursts

## Merging Together



Crawford et al. 2016


### Blue fraction



Butchler & Oemler 1984

Haines et al. 2009

## Substructure




Dressler-Shectman statistic is the classic test for substructure:

$$\delta^{2} = \frac{N_{obj}}{\sigma^{2}} [(\bar{v}_{local} - \bar{v})^{2} + ((\sigma_{local} - \sigma)^{2}].$$


Calculated as the offset from the cluster mean for the 10 nearest neighbors

## **Different Scales**




#### Overview

- Connecting populations
- Luminous Compact Blue Galaxies are triggered in galaxy clusters
- LCBGs have similar properties as dE
- Fate of LCBGs



#### Other Populations



## Magnitude-size

